

# Ethylene Insensitivity Modulates Ozone-Induced Cell Death in Birch<sup>1</sup>

Jorma Vahala, Raili Ruonala, Markku Keinänen<sup>2</sup>, Hannele Tuominen<sup>3</sup>, and Jaakko Kangasjärvi\*

Institute of Biotechnology and Department of Biosciences, University of Helsinki, POB 56 (Viikinkaari 9), FIN-00014 Helsinki, Finland (J.V., R.R., M.K., H.T., J.K.); Department of Applied Biology, University of Helsinki, POB 27 (Latokartanonkaari 5–7), FIN-00014 Helsinki, Finland (J.V.); and Laboratory of Plant Physiology and Molecular Biology, Department of Biology, University of Turku, FIN-20014 Turku, Finland (J.K.)

We have used genotypic variation in birch (*Betula pendula* Roth) to investigate the roles of ozone (O<sub>3</sub>)-induced ethylene (ET), jasmonic acid, and salicylic acid in the regulation of tissue tolerance to O<sub>3</sub>. Of these hormones, ET evolution correlated best with O<sub>3</sub>-induced cell death. Disruption of ET perception by transformation of birch with the dominant negative mutant allele *etr1-1* of the Arabidopsis ET receptor gene *ETR1* or blocking of ET perception with 1-methylcyclopropene reduced but did not completely prevent the O<sub>3</sub>-induced cell death, when inhibition of ET biosynthesis with aminooxyacetic acid completely abolished O<sub>3</sub> lesion formation. This suggests the presence of an ET-signaling-independent but ET biosynthesis-dependent component in the ET-mediated stimulation of cell death in O<sub>3</sub>-exposed birch. Functional ET signaling was required for the O<sub>3</sub> induction of the gene encoding  $\beta$ -cyanoalanine synthase, which catalyzes detoxification of the cyanide formed during ET biosynthesis. The results suggest that functional ET signaling is required to protect birch from the O<sub>3</sub>-induced cell death and that a decrease in ET sensitivity together with a simultaneous, high ET biosynthesis can potentially cause cell death through a deficient detoxification of cyanide.

The concentration of tropospheric ozone (O<sub>3</sub>) has increased during the past decades due to human activities, and it has been estimated that in year 2100, 50% of global forest area will be exposed to potentially phytotoxic O<sub>3</sub> concentrations (Fowler et al., 1999). In the leaves of O<sub>3</sub>-sensitive plants, symptoms of high O<sub>3</sub> are observed as rapid lesion formation. Traditionally, formation of reactive oxygen species (ROS), such as superoxide (O<sub>2</sub><sup>•−</sup>) and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) from the degradation of O<sub>3</sub> in the apoplast has been thought to alter the integrity of the plasma membrane and thus the integrity of the cell (Laisk et al., 1989; Heath, 1994). However, O<sub>3</sub> also induces an active and controlled apoptotic oxidative burst, the production of O<sub>2</sub><sup>•−</sup> and H<sub>2</sub>O<sub>2</sub> in the leaves affected, which may initiate programmed cell death analogous to that induced by ROS in an incompatible plant-pathogen interaction (Schraudner et al., 1998;

Pellinen et al., 1999, 2002; Overmyer et al., 2000; Moeder et al., 2002; Wohlgenuth et al., 2002).

Activation of ethylene (ET) biosynthesis by induction of the genes encoding 1-aminocyclopropane-1-carboxylate synthase (ACS) is one of the fastest and most obvious biochemical responses to O<sub>3</sub> and has been mechanistically linked to the regulation of O<sub>3</sub> lesion formation (Schlagnhauer et al., 1997; Tuominen et al., 1997; Vahala et al., 1998; Overmyer et al., 2000; Moeder et al., 2002). ET is perceived by a two-component His kinase receptor family (Chang et al., 1993a; Hua et al., 1995, 1998; Sakai et al., 1998). The ET receptor can be pharmacologically blocked with a competitive inhibitor of ET action, norbornadiene, or with 1-methylcyclopropene (MCP; Serek et al., 1995), which prevents the binding of ET to the receptor. The known dominant ET receptor mutations also prevent ET perception and thus cause ET insensitivity (Hall et al., 1999). This has been used to create stable ET insensitivity in petunia (*Petunia hybrida*), tomato (*Lycopersicon esculentum*), and tobacco (*Nicotiana tabacum*) by transforming them with the Arabidopsis *etr1-1* mutant allele (Wilkinson et al., 1997; Knoester et al., 1998).

Inhibition of ET biosynthesis with inhibitors of ACS has significantly reduced O<sub>3</sub>-induced lesion formation in the leaves of O<sub>3</sub>-exposed plants (Mehlhorn and Wellburn, 1987; Mehlhorn et al., 1991; Schlagnhauer et al., 1995; Wenzel et al., 1995; Tuominen et al., 1997; Moeder et al., 2002). Similarly, defective ET signaling in the ET-insensitive *ein2* mutant (Overmyer et al., 2000) or inhibition of ET per-

<sup>1</sup> This work was supported by the Maj and Tor Nessling Foundation, by the Finnish Society of Forest Sciences, by the Leo and Regina Wainstein's Foundation, and by the Academy of Finland, Centre of Excellence Program 2000–2005. M.K. was supported by Academy of Finland Postdoctoral grant no. 48640.

<sup>2</sup> Present address: Department of Biology, University of Joensuu, POB 111, FIN-80101 Joensuu, Finland.

<sup>3</sup> Present address: Umeå Plant Science Center, Umeå University, 90187 Umeå, Sweden.

\* Corresponding author; e-mail jaakko.kangasjarvi@helsinki.fi; fax 358–2–333–5549.

Article, publication date, and citation information can be found at [www.plantphysiol.org/cgi/doi/10.1104/pp.102.018887](http://www.plantphysiol.org/cgi/doi/10.1104/pp.102.018887).

ception with exogenous norbornadiene reduced O<sub>3</sub>-induced cell death in tomato (Bae et al., 1996; Moeder et al., 2002) and the spread of O<sub>3</sub>-induced programmed cell death in an O<sub>3</sub>-sensitive *Arabidopsis radical induced cell death1 (rcd1)* mutant (Overmyer et al., 2000). Furthermore, in tomato, both ET biosynthesis and the subsequent H<sub>2</sub>O<sub>2</sub> accumulation, which was required for the O<sub>3</sub> lesion formation, were highly colocalized, and inhibition of ET synthesis or perception also significantly reduced H<sub>2</sub>O<sub>2</sub> accumulation and O<sub>3</sub> lesion formation (Moeder et al., 2002). Taken together, in tomato and *Arabidopsis*, both ET synthesis and signaling are required in the processes that result in O<sub>3</sub> lesion formation.

In addition to ET, jasmonic acid (JA), and salicylic acid (SA) are involved in the regulation of oxidative stress responses in plants. JA is commonly regarded as a protective compound (Creelman and Mullet, 1995; Berger, 2001) and may have an essential role in the control of proper lesion containment upon pathogen and O<sub>3</sub> challenge (Thomma et al., 1998; Overmyer et al., 2000; Berger, 2001). SA is a central component in plant pathogen defense. It is involved in the regulation of the oxidative burst, cell death in hypersensitive response, defense signaling, and systemic acquired resistance (Godiard et al., 1994; Staskawicz et al., 1995; Draper, 1997; Shirasu et al., 1997; Ciardi et al., 2000; Dangl and Jones, 2001). SA has a dual role in the control of plant O<sub>3</sub> responses; SA promotes the oxidative cell death during a short-term O<sub>3</sub> challenge but is also required for the up-regulation of defenses during a long-term oxidative challenge (Rao and Davis, 1999, 2001).

The mechanisms of stress resistance and acclimation in trees are not well understood; for example, there are only a few studies on the hormonal interactions during oxidative stress in trees. In hybrid poplar (*Populus maximowiczii* × *P. trichocarpa*), sensitivity to acute high O<sub>3</sub> correlated with deficiencies in both JA- and SA-signaling pathways (Koch et al., 1998, 2000). However, the role of ET was unexplored. We have explored the role of ET under low chronic O<sub>3</sub> in the accompanying study (Vahala et al., 2003), which shows that ET also modifies O<sub>3</sub> sensitivity in hybrid aspen. In the experiments reported here, we have characterized the involvement of all three signaling pathways, ET, JA, and SA, in cell death induced by a short exposure to high O<sub>3</sub> concentration in silver birch (*Betula pendula* Roth) genotypes differing in their O<sub>3</sub> sensitivity. The role of ET was elucidated in detail using transgenic ET-insensitive birch lines and inhibitors. We show that ET insensitivity in birch reduced but did not completely eliminate O<sub>3</sub>-induced cell death when inhibition of ET biosynthesis abolished the lesion formation. This suggests the presence of an ET-signaling-independent but ET biosynthesis-dependent component in the ET-mediated stimulation of cell death in O<sub>3</sub>-exposed birch.

## RESULTS

### Kinetics of O<sub>3</sub>-Induced ET, JA, and SA Production among Three Wild-Type Birch Clones

Three birch clones, the O<sub>3</sub>-tolerant E9702, the moderately sensitive K1898 and the sensitive V5818 were selected from 17 genotypes differing in their O<sub>3</sub> sensitivity. One-year-old copies of the clones (ramets) were exposed to 200 nL L<sup>-1</sup> O<sub>3</sub> for 8 h. Over the whole experimental period, the clones did not differ statistically significantly in O<sub>3</sub>-induced ion leakage but displayed a differential pattern in O<sub>3</sub>-induced ion leakage (Fig. 1A).


ET evolution was highest in the O<sub>3</sub>-sensitive V5818 and lowest in the tolerant E9702. ET evolution was maximal at 4 to 8 h and correlated with ion leakage in all three clones (Table I). All of the changes in ET evolution were caused by O<sub>3</sub> because ET production did not change in air-grown plants throughout the experiments (data not shown). O<sub>3</sub> induced a significant increase in the accumulation of JA at 8 h in the O<sub>3</sub>-sensitive clone V5818, whereas only a slight increase was observed in K1898 and no increase was observed in the O<sub>3</sub>-tolerant E9702 (Fig. 1C). SA accumulated in significant amounts only 24 h after the onset of O<sub>3</sub> exposure, especially in the O<sub>3</sub> tolerant clone E9702. Clone K1898 accumulated intermediate levels of SA, whereas no SA accumulation was evident in the O<sub>3</sub>-sensitive V5818 (Fig. 1D). O<sub>3</sub>-induced ion leakage, JA and SA concentrations, and ET evolution increased more rapidly in clone K1898 than in the two other clones.

### Kinetics of Transcript Accumulation for Genes of ET Biosynthesis Correlates with O<sub>3</sub> Tolerance

Three birch ACS cDNAs were isolated to study whether the expression kinetics of ACS genes differed among the clones. In E9702, transcript levels for *BP-ACS1* (AY120897) and *BP-ACS2* (AY120898) reached the maximum at 4 h, whereas in V5818 and K1898, the maximum was reached at 8 h (Fig. 2, A and B). In V5818, the steady-state transcript levels for *BP-ACS1* and *BP-ACS2* were approximately 140- and 30-fold higher, respectively, than the initial level at 8 h, whereas in E9702, transcripts for *BP-ACS2* were almost undetectable at 8 h. *BP-ACS3* (AY120899) expression was not affected by O<sub>3</sub> (data not shown). In all clones, *BP-ACO1* (l-aminocyclopropane-l-carboxylate oxygenase; Y10749) transcript levels decreased between 0.5 and 1 h, increased back to the initial level at 2 h, and thereafter again decreased (Fig. 2C). The increase in transcript accumulation for *BP-ACO2* (AY154649) was among the fastest responses to O<sub>3</sub> and attained the maximum in clone E9702 at 2 h, and in clones K1898 and V5818 at 8 h (Fig. 2D).

### Generation of Transgenic, ET-Insensitive Birch Lines

Of the hormones investigated, ET evolution had the highest correlation with O<sub>3</sub>-induced cell death in all three clones (Table I). The role of ET in O<sub>3</sub>-induced cell death was studied further with inhibitors of ACS and ET perception and by generating in five different genetic backgrounds several transgenic birch lines that carry the dominant *etr1-1* mutant allele of the



**Figure 1.** Ion leakage, ET evolution, and JA and SA concentrations in three 1-year-old wild-type birch clones, V5818, K1898, and E9702, in response to O<sub>3</sub>. Clones were subjected to 200 nL L<sup>-1</sup> O<sub>3</sub> for 8 h, and leaves were harvested for ion leakage and hormone determinations after the beginning of the exposure at the times indicated. Black bar indicates the duration of O<sub>3</sub> fumigation. Measurements were conducted from three independent ramets at each time point/clone. Two-way ANOVA (clone and time as factors) followed by Tukey's honestly significant difference (HSD) mean-separation test was used to detect statistically significant differences between the clones. A, Ion leakage ( $P = 0.163$ ); B, ET evolution ( $P = 0.074$ ); C, free JA concentrations (ANOVA,  $P < 0.0005$ ; E9702 had lower JA accumulation than V5818 and K1898, Tukey's HSD mean-separation test  $P < 0.0005$  and  $P = 0.001$ , respectively); D, free SA concentrations (ANOVA,  $P < 0.0005$ ; K1898 had higher SA accumulation than V5818 and E9702, Tukey's HSD mean-separation test  $P < 0.0005$  and  $P = 0.006$ , respectively).

**Table I.** Correlation analysis of ion leakage, ET, JA, and SA in one-year-old clones V5818, K1898, and E9702 in response to O<sub>3</sub>.

The clones were subjected to 200 nL L<sup>-1</sup> of O<sub>3</sub> for 8 h, and leaves were harvested for ion leakage and hormone determinations at 0, 0.5, 1, 2, 4, 8, and 24 h from the beginning of the O<sub>3</sub>-exposure. Correlation analysis was conducted with the nonparametric Spearman's  $\rho$ . The statistical differences were considered significant at the level of  $P < 0.05$ . \*\* $P < 0.01$ ; \* $P < 0.05$ .

Clone	Hormone	Ion Leakage	ET	JA
V5818	ET	0.643**		
	JA	0.552**	0.582**	
	SA	0.173	0.121	0.392
K1898	ET	0.477*		
	JA	0.512*	0.376	
	SA	-0.004	-0.016	0.371
E9702	ET	0.469*		
	JA	-0.053	-0.022	
	SA	0.110	0.438	-0.001


Arabidopsis ET receptor gene *ETR1*. The degree of ET insensitivity in the transgenic lines was assessed by ET-induced leaf abscission and induction of an ET-regulated gene. After a 3-d exposure to ET, most of the wild type but none of the transgenic trees had freely abscised their leaves (Fig. 3D). When all of the remaining leaves were pulled with a rather strong force, a greatly attenuated leaf abscission was evident in 17 of the 23 independent transgenic lines examined (Fig. 3A). Southern analysis with the Arabidopsis *ETR1* revealed that the number of *etr1-1* inserts varied between the transgenic lines. For example, line 35 had two *etr1-1* inserts, and line 86 had one (Fig. 3B).

ET insensitivity was also visible as impaired ET induction of gene expression in the transgenic lines. A birch mitochondrial phosphate translocator gene (*BP-MPT1* [Y08499]; Kiiskinen et al., 1997) was strongly induced by ET in the wild-type birch clones but not in the transgenic lines. This was tested in five wild-type clones and 12 independent transgenic lines with similar results; the results for two of the transgenic lines (35 and 86) and the corresponding wild-type clone V5834 are shown in Figure 3C. We selected line 35, denoted hereafter as *BPetr1-1-35*, for further studies on the role of ET perception in O<sub>3</sub>-induced cell death. The O<sub>3</sub> sensitivity of V5834 is similar to clone V5818 used in the other experiments (J. Vahala, H. Tuominen, and J. Kangasjärvi, unpublished data).

### ET Insensitivity Reduces But Does Not Prevent O<sub>3</sub>-Induced Cell Death

The ET-insensitive *BPetr1-1-35* had less O<sub>3</sub> damage (two-way ANOVA; genotype and time as factors;  $P = 0.001$ ; Fig. 4A) but similar ET evolution (two-way ANOVA;  $P = 0.944$ ; Fig. 4B) when compared with the wild-type clone V5834. The remaining O<sub>3</sub> lesions were mainly localized near the veins. Because the





**Figure 2.** Relative increases in transcript abundance for the enzymes of ET biosynthesis in three 1-year-old wild-type birch clones, V5818, K1898, and E9702, in response to O<sub>3</sub>. Clones were subjected to 200 nL L<sup>-1</sup> O<sub>3</sub> for 8 h, and leaves for RNA isolation were harvested at the times indicated. Hybridization signals of each transcript are indicated as -fold increase of the initial level at 0 h. Black bars indicate the duration of O<sub>3</sub> fumigation. A, mRNA levels of *BP-ACS1*; B, mRNA levels of *BP-ACS2*; C, mRNA levels of *BP-ACO1*; and D, mRNA levels of *BP-ACO2*.

transgenic lines are insensitive to ET throughout their development and thus might lack some ET-dependent constitutive defenses, we elucidated the significance of ET signaling in the wild-type birch during a short-term O<sub>3</sub> exposure by blocking of ET perception with MCP only during the exposure. Similarly, as in the transgenic *BPetr1-1-35*, inhibition of ET perception in the wild-type clones partially reduced, but did not fully prevent visible O<sub>3</sub> lesion formation in the leaves of 1-year-old V5818 trees (Fig. 5A). Likewise, blocking of ET perception reduced the O<sub>3</sub>-induced visible cell death in 3-month-old leaves of clone V5818 (Fig. 5B).

To demonstrate that the MCP that could potentially be remaining in the cells did not cause any unexpected effects (for example, by possibly reacting


with O<sub>3</sub> during the exposure), we treated two wild-type birch clones, V5818 (Fig. 5C) and K1898 (not shown), with MCP immediately after a 6-h O<sub>3</sub> pulse. The blocking of ET perception with MCP after the O<sub>3</sub> treatment, again, only reduced cell death in both clones. This also indicates that ET is required for the O<sub>3</sub> lesion formation after the actual contact of O<sub>3</sub> with the apoplastic structures. In the transgenic *BPetr1-1-35*, O<sub>3</sub> damage was not reduced any further by a MCP treatment, whereas in the wild-type V5834, the extent of visual O<sub>3</sub> lesions was reduced by 32% (data not shown). Thus, the transformation of the genomic clone of the Arabidopsis *etr1-1* allele caused such a strong ET insensitivity in birch that it could not be further increased by exogenous MCP.

### Inhibition of ET Biosynthesis Prevents O<sub>3</sub>-Induced Cell Death

The ACS inhibitor AOA was used to determine how prevention of ET biosynthesis affects O<sub>3</sub> lesion formation in wild-type clone V5834 and the transgenic line *BPetr1-1-35*. AOA abolished ET production (three-way ANOVA; genotype, time and treatment as factors; Tukey's HSD mean-separation test;  $P < 0.0005$ ) and prevented the O<sub>3</sub>-dependent cell death completely in both wild-type and transgenic trees (Tukey's HSD mean-separation test;  $P = 0.001$ ; Fig. 4, C and D). Similarly, AOA treatment of either 3-month-old (data not shown) or 1-year-old V5818 ramets abolished O<sub>3</sub>-induced formation of visible lesions (Fig. 5D) and eliminated O<sub>3</sub>-induced ion leakage (three-way ANOVA; genotype, time and treatment as factors;  $P < 0.0005$ ) and ET evolution completely (three-way ANOVA;  $P < 0.0005$ ; data not shown). For an unknown reason, treatment with Tween 20 alone also reduced slightly the O<sub>3</sub>-induced ion leakage and ET evolution. Measured in wild-type clone V5818, the stomatal conductance in AOA-treated leaves was approximately 2-fold higher than either in O<sub>3</sub> or O<sub>3</sub>/MCP-treated leaves (data not shown). Thus, differential O<sub>3</sub> lesion formation under MCP and AOA treatments was not based on higher entry of O<sub>3</sub> to the leaf.

### The O<sub>3</sub>-Induced Activation of $\beta$ -Cyano-Ala Synthase ( $\beta$ -CAS) Is Dependent on ET

The results above suggest that the ET-dependent cell death is differentially affected by ET synthesis and signaling and prompts the question of what is the basis of cell death in the O<sub>3</sub>-exposed ET-insensitive plants that can be prevented by inhibition of ET biosynthesis. The ACO-catalyzed oxidation of 1-aminocyclopropane-1-carboxylic acid (ACC) to ET produces an equimolar amount of cyanide (HCN), which is normally rapidly detoxified to  $\beta$ -cyano-Ala by  $\beta$ -CAS (Akopyan et al., 1975; Wurtele et al., 1985; Yip and Yang, 1988). We examined whether the birch



**Figure 3.** Evidence for ET insensitivity in the transgenic birch lines carrying Arabidopsis ET receptor gene with the dominant *etr1-1* mutation. A, Leaf abscission in response to 50  $\mu\text{L L}^{-1}$  ET for 3 d in wild-type clones 4212, JR1/4, K1316, K1659, and V5834 and in 17 transgenic birch lines. Error bars indicate  $\pm\text{SE}$  ( $n = 6-11$ ). Leaf abscission in each clone was assessed by pulling with similar force every leaf of the trees exposed to ET. B, Southern analysis for the presence of the Arabidopsis *etr1-1* gene in the wild-type clone V5834 and in transgenic birch lines 35 and 86. Birch genomic DNA was digested with *KpnI* and was hybridized with an Arabidopsis *ETR1*-specific probe. C, Accumulation of birch mitochondrial phosphate translocator (*BP-MPT1*) transcript in the wild-type clone V5834 and in the transgenic birch lines 35 and 86 in response to 50  $\mu\text{L L}^{-1}$  ET. Black bar indicates the duration of ET treatment. D, Leaf abscission in response to a 3-d ET treatment in the wild type (left) and *etr1-1* transformed transgenic line (right) of birch.

$\beta$ -CAS gene (AY154650) is ET regulated. Results in Figure 6A show that when in the wild-type clone V5834  $\beta$ -CAS transcript accumulation increased over 16-fold by exposure to ET, in the ET-insensitive *BPetr1-1-35*  $\beta$ -CAS was not induced. In response to O<sub>3</sub>,  $\beta$ -CAS transcript accumulation was also lower in *BPetr1-1-35* than in the wild-type V5834 (Fig. 6B), whereas ET evolution was equal (Fig. 4B). Similarly, blocking of ET receptors with MCP reduced  $\beta$ -CAS transcript accumulation in response to O<sub>3</sub>, but ET evolution was not affected (Fig. 6C). Thus, it can be concluded that functional ET signaling is required for the O<sub>3</sub>-induced activation of  $\beta$ -CAS in birch and that as a result of decreased ET sensitivity, HCN detoxification can be compromised.


## DISCUSSION

### O<sub>3</sub>-Induced Cell Death Correlates with ET and JA Accumulation

According to the recent results in several studies (Rao and Davis, 2001), plant responses to high concentrations of O<sub>3</sub> share common features with pathogen attack due to the function of ROS as signaling molecules. In this study, we employed high O<sub>3</sub> concentrations as a tool to probe the interactions of hor-

monal responses and the high level of ROS in birch. The results of these experiments address more the interactions and roles of SA, JA, and ET with ROS in the regulation of ROS-dependent cell death, and thus are not directly applicable to describe or represent plant responses to O<sub>3</sub> in natural environments. The latter we have evaluated in the accompanying paper (Vahala et al., 2003; this issue), where the roles of same hormones in response to chronic O<sub>3</sub> concentrations have been studied in hybrid aspen (*Populus tremula*  $\times$  *P. tremuloides*) under more realistic levels of O<sub>3</sub> found in natural environments.


The timing and magnitude of ET evolution correlated with the extent of cell death in the wild-type birch clones. As could be expected, blocking of ET biosynthesis in the O<sub>3</sub>-sensitive clone V5818 reduced O<sub>3</sub>-induced cell death significantly, indicating the requirement for ET. The O<sub>3</sub>-sensitive clone V5818 displayed high O<sub>3</sub>-induced ET accumulation without the late SA accumulation, whereas the O<sub>3</sub>-tolerant clone E9702 had low O<sub>3</sub>-induced ET and highly induced late SA accumulation. This suggests that the early high ET production may antagonize the late SA accumulation, or vice versa, increased SA production may down-regulate ET accumulation and thus prevent the ET-dependent cell death.



**Figure 4.** O<sub>3</sub>-induced ion leakage, ET evolution, and the effect of blocking of ET biosynthesis in 3-month-old birch clone V5834 and in the transgenic line *BPetr1-1-35* carrying *Arabidopsis* ET receptor gene with the dominant *etr1-1* mutation. Ramets were sprayed either with 0.05% (v/v) Tween 20 or with 1.5 mM aminooxyacetic acid (AOA) in 0.05% (v/v) Tween 20 to block ET biosynthesis and were exposed to 200 nL L<sup>-1</sup> O<sub>3</sub> for 6 h. Leaves were harvested for ET and ion leakage determinations at 0, 6, 8, and 24 h. Error bars indicate  $\pm$ SE ( $n = 3$ ). Black bars indicate the duration of O<sub>3</sub> exposure. A, Ion leakage in the wild-type clone V5834 and in the transgenic line *BPetr1-1-35*. B, ET evolution in the wild-type clone V5834 and in the transgenic line *BPetr1-1-35*. C, Ion leakage in the wild-type clone V5834 and in the transgenic line *BPetr1-1-35* with and without the AOA treatment. D, ET evolution in the wild-type clone V5834 and in the transgenic line *BPetr1-1-35* with and without the AOA treatment.


Similarly, the basal SA level of the ET-insensitive *Arabidopsis* mutant *ein2* was higher than in wild type, suggesting that ET can inhibit SA biosynthesis (H. Tuominen, K. Overmyer, M. Keinänen, and J. Kangasjärvi, unpublished data). The same interaction works also in the opposite direction; O<sub>3</sub>-induced, EIN2-dependent gene expression was suppressed in the SA-deficient NahG plants. These results suggest similar regulation and interaction between ET and SA during oxidative stress in both *Arabidopsis* and birch. The role of JA is, however, more complicated. In birch, both ET and JA accumulation correlated with O<sub>3</sub>-induced cell death in the O<sub>3</sub>-sensitive clones V5818 and K1898, whereas in the O<sub>3</sub>-tolerant clone E9702, JA concentration did not increase. Similarly, in the O<sub>3</sub>-sensitive *Arabidopsis* mutants *rcd1* and *jar1*, high JA accumulation was evident. It has been shown that JA is involved in lesion containment (Overmyer et al., 2000), which at first may seem contradictory to the results of JA accumulation in the sensitive clones and not in the tolerant ones. However, because the biosynthesis of JA appears to be

limited by substrate availability (Laudert et al., 2000; Ziegler et al., 2001), it is also possible that the accumulation of JA is a consequence of the ET-dependent cell death: Substrate for JA synthesis ( $\alpha$ -linolenic acid or 13-(S)-hydroperoxylinolenic acid) could be released from the dying cells, thus resulting in in-



**Figure 5.** Visual damage in response to O<sub>3</sub> in birch after the inhibition of ET perception or biosynthesis. ET perception was blocked with 300 nL L<sup>-1</sup> MCP and ET biosynthesis with 1 mM AOA. Ramets were exposed to 200 nL L<sup>-1</sup> O<sub>3</sub> for 6 h. Photographs were taken 48 h after the beginning of the O<sub>3</sub> exposure. A, An O<sub>3</sub>-treated leaf of the 1-year-old clone V5818 (left) and a leaf pretreated with MCP and then with O<sub>3</sub> (right). B, A detail of an O<sub>3</sub>-treated leaf of the 3-month-old clone V5818 (left) and a leaf pretreated with MCP and then with O<sub>3</sub> (right). C, A detail of an O<sub>3</sub>-treated leaf of the 3-month-old clone V5818 (left) and a leaf treated with MCP (right) after the O<sub>3</sub> pulse. D, An O<sub>3</sub>-treated leaf of the 1-year-old clone V5818 (left) and a leaf treated with AOA and O<sub>3</sub> (right).





**Figure 6.** Accumulation of  $\beta$ -CAS transcript in response to ET or O<sub>3</sub>. Birch clone V5834, the transgenic *BPetr1-1-35* line carrying the Arabidopsis ET receptor gene with the dominant *etr1-1* mutation, and clone V5818 with blocked ET perception were exposed to O<sub>3</sub> or ET, and the transcript levels of  $\beta$ -CAS were analyzed. Ramets were subjected to 50  $\mu\text{L L}^{-1}$  ET for 24 h or 200 nL L<sup>-1</sup> O<sub>3</sub> for 6 h. ET perception was blocked with 300 nL L<sup>-1</sup> MCP. Black bars indicate the duration of treatments. A,  $\beta$ -CAS mRNA accumulation in response to ET treatment in the wild-type clone V5834 and in the transgenic *BPetr1-1-35* line. B,  $\beta$ -CAS mRNA accumulation in response to O<sub>3</sub> in the wild-type clone V5834 and in the transgenic line *BPetr1-1-35*. C  $\beta$ -CAS mRNA accumulation in response to O<sub>3</sub> in the control and MCP-treated wild-type clone V5818. Inset, ET evolution in the control and MCP-treated ramets in response to O<sub>3</sub>.

creased JA synthesis, which then halts the ET-dependent lesion propagation.

#### A Dual Role for ET in O<sub>3</sub> Responses

ET can have opposite roles in O<sub>3</sub>-exposed plants. Several reports have shown an essential pro-death role for ET in O<sub>3</sub>-exposed plants. However, depending on the temporal pattern of biosynthesis ET may also have a pro-survival role. In mung bean (*Vigna radiata*) and pea (*Pisum sativum*), pretreatment of plants with ET before O<sub>3</sub> exposure promoted O<sub>3</sub> tol-

erance (Mehlhorn, 1990). Involvement of ET-dependent protection from oxidative stress may require a sufficient level of rapid but transient ET induction, however, at the intensity where it does not exceed the rate that induces prolonged spreading cell death. In the O<sub>3</sub>-sensitive clone V5818, timing and degree of the late ET accumulation was most likely beyond the limit to provide protection against the O<sub>3</sub> challenge. This is also supported by the fact that blocking of ET perception after the O<sub>3</sub> exposure decreased O<sub>3</sub> damage, which indicates that the ET action for the promotion of cell death is required late. Although the early transcript accumulation of *BP-ACS1* was evident in all birch clones in response to O<sub>3</sub>, *BP-ACS2* transcript levels increased considerably only in the O<sub>3</sub>-sensitive clone V5818 at 8 h. Therefore, in these three wild-type clones, differential ACS transcript accumulation and ET production suggest a dual role for ET. Depending on the magnitude of synthesis and its temporal pattern, ET can serve as a mediator of either survival or cell death. The results presented here suggest that in the O<sub>3</sub>-sensitive birch clone ET was required to promote cell death at the time of lesion development and that the requirement of ET, for example, for the induction of antioxidant defenses had less significance.

#### Transformation of Arabidopsis *etr1-1* Allele Causes Strong ET Insensitivity in Birch

The dominant Arabidopsis *etr1-1* mutation causes ET insensitivity (Chang et al., 1993a). When the genomic clone of Arabidopsis *etr1-1* was transformed to tobacco, it caused ET insensitivity in the transgenic plants and strongly reduced basic pathogenesis-related-gene expression and non-host resistance against *Pythium* spp. (Knoester et al., 1998). Here, we show that transformation of birch with the genomic copy of the Arabidopsis *etr1-1* mutant allele under its own promoter resulted in ET insensitivity in birch. Furthermore, the ET insensitivity was strong, which is demonstrated by the fact that MCP treatment of the transgenic trees did not decrease the ET-dependent O<sub>3</sub> lesion formation any further. Therefore, regulation of *ETR1* is most likely conserved among various plant species because apparently birch transcription factors recognized the Arabidopsis *ETR1* promoter and regulated its expression properly.

#### O<sub>3</sub> Lesions in ET-Insensitive Plants May Be Due to Defective HCN Removal

Insensitivity to ET provided only a partial protection against O<sub>3</sub>, when the prevention of ET biosynthesis in the ET-insensitive transgenic birch blocked cell death completely. This suggests that ET modifies O<sub>3</sub>-induced ET-dependent cell death downstream of ACS and both upstream and downstream of ET receptors. When ACC is oxidized to ET, stoichiometric amounts of ET and HCN are produced. Under nor-

mal circumstances, this HCN is efficiently detoxified by  $\beta$ -CAS. However, it has been suggested (Grossmann, 1996) that under specific conditions, strongly stimulated ET biosynthesis may result in necrotic cell death due to insufficient HCN detoxification. Infection of tobacco with tobacco mosaic virus caused elevated HCN and ACC levels, concomitant with a decrease in  $\beta$ -CAS activity (Siefert et al., 1995), and HCN accumulation was suggested to contribute to the cell death. Our results indicate that the birch  $\beta$ -CAS gene is ET-regulated. The  $\beta$ -CAS transcript accumulation was also considerably lower in response to  $O_3$  when ET perception was disrupted. However, this was still accompanied by equal  $O_3$ -induced ET accumulation when compared with the wild-type birch, thus ACO was still producing HCN at an equal rate as ET. Hence, the HCN formed could be a likely candidate to mediate the  $O_3$ -induced cell death.

HCN may also relate to the ROS formation in the mitochondria: It has been shown that the HCN-resistant respiration decreased mitochondrial ROS formation in cultured tobacco cells via the alternative oxidase (AOX; Maxwell et al., 1999). It has also been shown with the Arabidopsis *etr1-1* mutant that AOX activation is ET dependent (Simons et al., 1999). The AOX gene is also induced by  $H_2O_2$  (Robson and Vanlerberghe, 2002). Pellinen et al. (1999, 2002) have shown that  $O_3$  induces  $H_2O_2$  accumulation in birch. This suggests that AOX was most likely induced as well. Thus, attenuated induction of the HCN-resistant respiration together with defective HCN removal in ET-insensitive plants may result in inhibition of the normal mitochondrial respiration by HCN and thus cause increased ROS production in the mitochondria, which could be involved in the regulation of cell death. When the subcellular compartmentalization of  $O_3$ -induced  $H_2O_2$  formation was studied in birch (Pellinen et al., 1999), increased ROS accumulation temporally coinciding with cell death was observed also in the mitochondria.

In this study, the defect in HCN detoxification may be considered an "artificial" effect caused by the strong ET insensitivity obtained with *etr1-1* transformation or MCP treatment. This strong ET insensitivity may be regarded as unlikely to be observed in wild-type plants. However, our results in Arabidopsis suggest that modulation of ET sensitivity takes place in  $O_3$ -exposed plants (H. Tuominen, K. Overmyer, M. Keinänen, and J. Kangasjärvi, unpublished data). Furthermore, it has been shown that in tomato, the genes encoding ET receptors are differentially induced by  $O_3$  (Moeder et al., 2002) and during pathogen infection (Ciardi et al., 2000). The increased synthesis of "fresh," unoccupied receptors has been proposed to decrease ET sensitivity and to be involved in the desensitization of plants to ET when the ET responses need to be shut down (Bleecker and Kende, 2000; Ciardi and Klee, 2001). In tomato, the

$O_3$ -induced ET synthesis, ET-dependent  $H_2O_2$  accumulation, and subsequent cell death were highly colocalized to the vicinity of the veins (Moeder et al., 2002). The ET-signaling-independent lesion formation in birch was also near the vascular system. Thus, it is completely possible that the simultaneous and extremely localized high ET biosynthesis in connection with decreased ET sensitivity caused by the induction of ET receptor genes can cause cell death through increased HCN formation and mitochondrial responses.

The deficient HCN removal in transgenic, ET-insensitive plants also has implications in other species that have been made insensitive to ET by transformation of mutant ET receptors; under conditions that cause highly elevated ET synthesis, sufficient HCN detoxification may not take place due to the ineffective induction of  $\beta$ -CAS.

## MATERIALS AND METHODS

### Plant Material and Growth Conditions

Three birch (*Betula pendula* Roth) genotypes (V5818, K1898, and E9702) and ET-insensitive birch lines were propagated by in vitro tissue culture as described previously (Lemmettyinen et al., 1998). Copies of the clones (ramets) were planted and grown in peat:sand:vermiculite (6:1:1, w/v), fertilized with nitrogen:phosphorus:potassium (11:4:25) under greenhouse conditions, and used for experiments during either the 1st (3 month old) or 2nd year of their growth. Ramets were transferred into growth chambers with the photoperiod of 22 h of light/2 h of dark, light intensity of  $300 \mu\text{mol m}^{-2} \text{s}^{-1}$  photosynthetically active radiation, temperature of  $20^\circ\text{C}/16^\circ\text{C}$  (light/dark), and relative humidity of 50%/70% (light/dark) and allowed to acclimate to the chamber conditions at least for 4 d before the  $O_3$ , ET, and inhibitor treatments.

ET-insensitive birch were created by genetic transformation with a construct (pCGN1547) carrying the Arabidopsis ET receptor gene *ETR1* with the dominant *etr1-1* mutation with its own promoter (Chang et al., 1993a). Different birch clones (4212, E9678, E9702, JR1/4, K1659, K1898, V5818, and V5834) were chosen for *Agrobacterium* sp.-mediated (C58C1 pGV2260) transformation as described (Lemmettyinen et al., 1998). Clones V5818, K1898, and E9702 used in the other experiments reported here could not be successfully transformed (data not shown).

### $O_3$ , Chemical, and ET Treatments

The  $O_3$  exposures were conducted in growth chambers.  $O_3$  was produced from pure  $O_2$  by electric discharge, and the delivery of  $O_3$  to the chambers was computer-controlled based on continuous measurements of  $O_3$  concentration inside the chamber with an ozone analyzer (Dasibi 1008-RS, Amko Systems Inc., Ontario, Canada) as described by Pellinen et al. (1999). Wild-type birch clones (V5818, K1898, and E9702) were exposed to  $200 \text{ nL L}^{-1} O_3$  for 8 h, and leaves were harvested at 0, 0.5, 1, 2, 4, 8, and 24 h. ET-insensitive birch ramets were exposed to  $200 \text{ nL L}^{-1} O_3$  for 6 h, and leaves were harvested at 0, 6, 8, and 24 h. Wild-type birch ramets of the same clone were used as controls.

ET receptors were blocked in the wild-type clones (V5818 and K1898) and ET-insensitive transgenic birch line *BPetr1-1-35* with 1-MCP (EthylBloc, Laboratorium Van der Sprong bv, Netherlands). The treatments were carried out for 12 h in a sealed growth chamber with  $300 \text{ nL L}^{-1}$  MCP according to manufacturer's instructions, before or after the  $O_3$  treatment. After each MCP pretreatment, the chamber was flow-through ventilated for 2 h before the  $O_3$  exposure to avoid any possible chemical reactions between  $O_3$  and MCP.

ET biosynthesis was blocked in the wild-type clone V5818 and transgenic birch line *BPetr1-1-35* with 1 mM or 1.5 mM AOA (Sigma-Aldrich) with 0.1% or 0.05% (v/v) Tween 20 (Fluka, Buchs, Switzerland), respectively, by spraying the abaxial side of the leaves. The control plants for AOA treat-



ment received 0.1% or 0.05% (v/v) Tween 20. Spraying was conducted 1.5 h before the onset of O<sub>3</sub> exposure and was repeated four times during the O<sub>3</sub> exposure.

ET treatment was conducted to test the ET sensitivity of the 3-month-old transgenic *etr1-1* clones by subjecting ramets to 50  $\mu\text{L L}^{-1}$  ET for 3 d in a growth chamber.

## ET, Ion Leakage, JA, and SA Determinations

The fully expanded leaves 1, 2, and 5 from the apex were pooled for ET evolution and ion leakage measurements from three ramets per clone per treatment. The ET and ion leakage determinations were carried out as described by Vahala et al. (2003).

The fully expanded leaves 3, 4, and 6 from the apex were used for JA and SA determinations at each time point. JA and SA were extracted and quantified with [1,2-<sup>13</sup>C]JA and [<sup>13</sup>C]SA as internal standards by gas chromatography-mass spectrometry as described by Baldwin et al. (1997), with the following modifications. The frozen tissues were ground in 2-mL microfuge tubes, spiked with an internal standard, extracted overnight with 1.5 mL of acetone:50 mM citric acid (7:3, v/v) at 4°C with vigorous shaking, and re-extracted for 15 min with 1 mL of the extraction solvent. After centrifugation (13,000 rpm for 8 min), the combined supernatants were divided in two aliquots (for free and conjugated compounds) and evaporated to an aqueous residue under vacuum at 30°C. For the analysis of conjugated SA, the sample was adjusted to 400  $\mu\text{L}$  with deionized water, acidified with 100  $\mu\text{L}$  of concentrated HCl (37%, w/v), and hydrolyzed for 1 h at 80°C. After hydrolysis, the aqueous solutions were extracted twice with 1 mL of diethyl ether and evaporated to dryness under vacuum at ambient temperature. The residues were redissolved in 1 mL of diethyl ether and loaded onto 100 mg of Supelclean LC-NH<sub>2</sub> SPE columns (Supelco, Bellefonte, PA). The columns were washed with 1.2 mL of chloroform:2-propanol (3:1, v/v), and the compounds were eluted with 1.5 mL of diethyl ether:formic acid (98:2, v/v). The eluates were evaporated to dryness under vacuum at ambient temperature, dissolved in 100  $\mu\text{L}$  of diethyl ether, derivatized with ethereal diazomethane, and reconstituted in 40  $\mu\text{L}$  of hexane.

The methyl esters of JA and SA were separated by gas chromatography (HP 6890, Agilent Technologies, Avondale, PA) on an Rtx-5MS column (30-m  $\times$  0.25-mm i.d., 0.25- $\mu\text{m}$  film thickness; Restek Corp., Bellefonte, PA) with a helium flow rate of 1 mL min<sup>-1</sup> and injector temperature of 250°C, and detected by mass spectrometry (HP 5973) in a single-ion-monitoring mode (*m/z* 224 and 226 for JA, *m/z* 152 and 153 for SA). The temperature program was as follows, 1.5 min at 55°C, 10°C min<sup>-1</sup> to 200°C, 20°C min<sup>-1</sup> to 300°C, and hold at 300°C for 7 min.

## cDNA Probes and Northern and Southern Analysis

Several degenerate oligonucleotide primers based on conserved amino acid domains were used to isolate cDNAs for the ET biosynthetic enzymes from birch. Reverse transcription of total RNA or poly(A<sup>+</sup>) mRNA was conducted with the avian myeloblastosis virus reverse transcriptase (Promega, Madison, WI), and the subsequent PCR was conducted with Dynazyme (Finnzymes, Espoo, Finland), *Taq*, or *Pfu* (Promega) DNA polymerases. To isolate birch ACS cDNA sequences, *OLE-4* and *OLE-5* upstream primers and *OLE-6* downstream primer were used as described (Botella et al., 1992).

Young leaf tissue collected from the apex of shoots was used for Southern analyses. Total DNA was extracted as described (Lodhi et al., 1994) and digested with *Kpn*I and *Eco*RI. For northern analysis, the fully expanded leaves 3, 4, and 6 from the apex were excised from three individuals per clone per treatment and frozen in liquid N<sub>2</sub> at each time point. Samples were stored at -70°C or -80°C until analyzed. The three independent samples were pooled, and total RNA was isolated as described (Chang et al., 1993b). Ten micrograms of total RNA was fractionated on 1% (w/v) formaldehyde agarose gels in MOPS buffer. RNA and DNA were capillary blotted overnight onto positively charged nylon membranes (Roche Diagnostics GmbH, Mannheim, Germany), and baked at 120°C for 30 min. [ $\alpha$ -<sup>32</sup>P]dCTP-labeled probes were prepared with High Prime random-priming labeling system (Roche Diagnostics) and purified on G-50 columns (Amersham Biosciences, Piscataway, NJ). Prehybridizations and hybridizations were carried out at 68°C in a solution containing 0.5 M NaHPO<sub>4</sub> and 7% (w/v) SDS. After

hybridization, the membranes were washed under high-stringency conditions as described (Church and Gilbert, 1984). Hybridization signals were quantified with a phosphor imager and an image analysis program (Bas 1500, Fujifilm, Tokyo) and normalized against the 18S rDNA hybridization signal.

## Photosynthesis Measurements

The net photosynthesis, stomatal conductance, and transpiration measurements were performed at 5 to 6 h after the onset of O<sub>3</sub> exposure with the LI-6400 photosynthesis system (LI-COR, Lincoln, NE). Measurements were performed on the third and fourth fully expanded leaves of three ramets per treatment. The mean value of two leaves measured from each individual ramet was calculated. All of the measurements were made under a 6400-02B red/blue LED light source at saturating light of 1,000  $\mu\text{mol m}^{-2} \text{s}^{-1}$  photosynthetically active radiation, temperature of 22°C, constant input of CO<sub>2</sub> (400  $\mu\text{L L}^{-1}$ ) from the minicartridges, constant air flow rate of 500  $\mu\text{mol s}^{-1}$ , and relative humidity of 30% to 40%.

## Statistical Analysis and Quantification of Visual Damage

ANOVA was used to detect significant differences among clones and treatments. All data were checked for normality and heterogeneity of variances. Hormone concentrations were log<sub>10</sub>(X) transformed and ion leakage arcsin transformed to meet the assumptions of ANOVA. Multiple comparisons of individual means and levels of factors were analyzed with a Tukey's HSD test, where appropriate. Correlations were analyzed with non-parametric Spearman's  $\rho$  test. Analyses were conducted with the SPSS v8.0 software package (SPSS Inc., Chicago). The O<sub>3</sub>-induced visual damage was estimated as a percentage of the leaf area that was damaged 48 h after the onset of the O<sub>3</sub> exposure. All fully emerged leaves were counted. Two people conducted the quantification independently.

## ACKNOWLEDGMENTS

We acknowledge Prof. Elliot Meyerowitz for providing the *etr1-1* genomic clone. We thank Kari Vaahomeri, Sari Eräluoto, Mirva Lehtinen, Airi Tauriainen, and Sari Möttönen for technical help. We acknowledge Kirk Overmyer for helpful discussions and Simo Harju and Mika Korva for the nursing of the birch ramets. We thank Prof. Ian Baldwin and Dr. Günter Brader for JA and SA standards, respectively.

Received December 9, 2002; returned for revision January 20, 2003; accepted February 19, 2003.

## LITERATURE CITED

- Akopyan T, Braunstein A, Goryachenkova E (1975) Beta-cyanoalanine synthase: purification and characterization. *Proc Natl Acad Sci USA* **72**: 1617-1621
- Bae GY, Nakajima N, Ishizuka K, Kondo N (1996) The role in ozone phytotoxicity of the evolution of ethylene upon induction of 1-aminocyclopropane-1-carboxylic acid synthase by ozone fumigation in tomato plants. *Plant Cell Physiol* **37**: 129-134
- Baldwin IT, Zhang Z-P, Diab N, Ohnmeiss TE, McCloud ES, Lynds GY, Schmelz EA (1997) Quantification, correlations and manipulations of wound-induced changes in jasmonic acid and nicotine in *Nicotiana sylvestris*. *Planta* **201**: 397-404
- Berger S (2001) Jasmonate-related mutants of *Arabidopsis* as tools for studying stress signals. *Planta* **214**: 497-504
- Bleecker AB, Kende H (2000) Ethylene: a gaseous signal molecule in plants. *Annu Rev Cell Dev Biol* **16**: 1-18
- Botella JR, Schlagnhauer CD, Arteca RN, Phillips AT (1992) Identification and characterization of three putative genes for 1-aminocyclopropane-1-carboxylate synthase from etiolated mung bean hypocotyl segments. *Plant Mol Biol* **18**: 793-797
- Chang C, Kwok SF, Bleecker AB, Meyerowitz EM (1993a) *Arabidopsis* ethylene-response gene *ETR1*: similarity of products to two-component regulators. *Science* **262**: 539-544

- Chang S, Puryear J, Cairney C (1993b) A simple and efficient method for isolating RNA from pine trees. *Plant Mol Biol Rep* **11**: 113–116
- Church GM, Gilbert W (1984) Genomic sequencing. *Proc Natl Acad Sci USA* **81**: 1991–1995
- Ciardi J, Klee H (2001) Regulation of ethylene-mediated responses at the level of receptor. *Ann Bot* **88**: 813–822
- Ciardi JA, Tieman DM, Lund ST, Jones JB, Stall RE, Klee HJ (2000) Response to *Xanthomonas campestris* pv. *vesicatoria* in tomato involves regulation of ethylene receptor gene expression. *Plant Physiol* **123**: 81–92
- Creelman RA, Mullet JE (1995) Jasmonic acid distribution and action in plants: regulation during development and response to biotic and abiotic stress. *Proc Natl Acad Sci USA* **92**: 4114–4119
- Dangl JL, Jones JDG (2001) Plant pathogens and integrated defence responses to infection. *Nature* **411**: 826–833
- Draper J (1997) Salicylate, superoxide synthesis and cell suicide in plant defence. *Trends Plant Sci* **2**: 162–165
- Fowler D, Cape NJ, Coyle M, Flechard C, Kuylensstierna J, Hicks K, Derwent D, Johnson C, Stevenson D (1999) The global exposure of forest to air pollutants. *Water Air Soil Pollut* **116**: 5–23
- Godiard L, Grant MR, Dietrich RA, Kiedrowski S, Dangl JL (1994) Perception and response in plant disease resistance. *Curr Opin Genet Dev* **4**: 662–671
- Grossmann K (1996) A role for cyanide, derived from ethylene biosynthesis, in the development of stress symptoms. *Physiol Plant* **97**: 772–775
- Hall AE, Chen QG, Findell JL, Schaller GE, Bleecker AB (1999) The relationship between ethylene binding and dominant insensitivity conferred by mutant forms of the ETR1 ethylene receptor. *Plant Physiol* **121**: 291–299
- Heath RL (1994) Possible mechanisms for the inhibition of photosynthesis by ozone. *Photosynth Res* **39**: 439–451
- Hua J, Chang C, Sun Q, Meyerowitz EM (1995) Ethylene-insensitivity conferred by *Arabidopsis* *ERS* gene. *Science* **269**: 1712–1714
- Hua J, Sakai H, Nourizadeh S, Chen QG, Bleecker AB, Ecker JR, Meyerowitz EM (1998) *EIN4* and *ERS2* are members of the putative ethylene receptor gene family in *Arabidopsis*. *Plant Cell* **10**: 1321–1332
- Kiiskinen M, Korhonen M, Kangasjärvi J (1997) Isolation and characterization of cDNA for a plant mitochondrial phosphate translocator (*Mpt1*): Ozone stress induces *Mpt1* mRNA accumulation in birch (*Betula pendula* Roth). *Plant Mol Biol* **35**: 271–279
- Knoester M, van Loon LC, van den Heuvel J, Hennig J, Bol JF, Linthorst JM (1998) Ethylene-insensitive tobacco lacks nonhost resistance against soil-borne fungi. *Proc Natl Acad Sci USA* **95**: 1933–1937
- Koch JR, Creelman RA, Eshita SM, Seskar M, Mullet JE, Davis KR (2000) Ozone sensitivity in hybrid poplar correlates with insensitivity to both salicylic acid and jasmonic acid: the role of programmed cell death in lesion formation. *Plant Physiol* **123**: 487–496
- Koch JR, Scherzer AJ, Eshita SM, Davis KR (1998) Ozone sensitivity in hybrid poplar is correlated with a lack of defense-gene activation. *Plant Physiol* **118**: 1243–1252
- Laik A, Kull O, Moldau H (1989) Ozone concentration in leaf intercellular air spaces is close to zero. *Plant Physiol* **90**: 1163–1167
- Laudert D, Schaller F, Weiler EW (2000) Transgenic *Nicotiana tabacum* and *Arabidopsis thaliana* plants overexpressing allene oxide synthase. *Planta* **211**: 163–165
- Lemmetyinen J, Keinonen-Mettälä K, Lännenpää M, von Weissenberg K, Sopanen T (1998) Activity of the CaMV 35S promoter in various parts of transgenic early flowering birch clones. *Plant Cell Rep* **18**: 243–248
- Lodhi MA, Ye G-N, Weeden NF, Reisch BI (1994) A simple and efficient method for DNA extraction from grapevine cultivars and *Vitis* species. *Plant Mol Biol Rep* **12**: 6–13
- Maxwell DP, Wang Y, McIntosh L (1999) The alternative oxidase lowers mitochondrial reactive oxygen production in plant cells. *Proc Natl Acad Sci USA* **96**: 8271–8276
- Mehlhorn H (1990) Ethylene-promoted ascorbate peroxidase activity protects plants against hydrogen peroxide, ozone and paraquat. *Plant Cell Environ* **13**: 971–976
- Mehlhorn H, O'Shea JM, Wellburn AR (1991) Atmospheric ozone interacts with stress ethylene formation by plants to cause visible plant injury. *J Exp Bot* **42**: 17–24
- Mehlhorn H, Wellburn AR (1987) Stress ethylene formation determines plant sensitivity to ozone. *Nature* **327**: 417–418
- Moeder W, Barry CS, Tauriainen AA, Betz C, Tuomainen J, Utriainen M, Grierson D, Sandermann H, Langebartels C, Kangasjärvi J (2002) Ethylene synthesis regulated by bi-phasic induction of 1-aminocyclopropane-1-carboxylic acid synthase and 1-aminocyclopropane-1-carboxylic acid oxidase genes is required for hydrogen peroxide accumulation and cell death in ozone-exposed tomato. *Plant Physiol* **130**: 1918–1926
- Overmyer K, Tuominen H, Kettunen R, Betz C, Langebartels C, Sandermann H Jr, Kangasjärvi J (2000) The ozone-sensitive *Arabidopsis rcd1* mutant reveals opposite roles for ethylene and jasmonate signaling pathways in regulating superoxide-dependent cell death. *Plant Cell* **12**: 1849–1862
- Pellinen R, Palva T, Kangasjärvi J (1999) Subcellular localization of ozone-induced hydrogen peroxide production in birch (*Betula pendula*) leaf cells. *Plant J* **20**: 349–356
- Pellinen RI, Korhonen M-S, Tauriainen AA, Palva ET, Kangasjärvi J (2002) H<sub>2</sub>O<sub>2</sub> activates cell death and defense gene-expression in birch (*Betula pendula*). *Plant Physiol* **130**: 549–560
- Rao MV, Davis KR (1999) Ozone-induced cell death occurs via two distinct mechanisms in *Arabidopsis*: the role of salicylic acid. *Plant J* **17**: 603–614
- Rao MV, Davis KR (2001) The physiology of ozone-induced cell death. *Planta* **213**: 682–690
- Robson CA, Vanlerberghe GC (2002) Transgenic plant cells lacking mitochondrial alternative oxidase have increased susceptibility to mitochondria-dependent and -independent pathways of programmed cell death. *Plant Physiol* **129**: 1908–1920
- Sakai H, Hua J, Chen QG, Chang C, Medrano LJ, Bleecker AB, Meyerowitz EM (1998) *ETR2* is an *ETR1*-like gene involved in ethylene signaling in *Arabidopsis*. *Proc Natl Acad Sci USA* **95**: 5812–5817
- Schlagghauser CD, Arteca RN, Pell EJ (1997) Sequential expression of two 1-aminocyclopropane-1-carboxylate synthase genes in response to biotic and abiotic stresses in potato (*Solanum tuberosum* L.) leaves. *Plant Mol Biol* **35**: 683–688
- Schlagghauser CD, Glick RE, Arteca RN, Pell EJ (1995) Molecular cloning of an ozone-induced 1-aminocyclopropane-1-carboxylate synthase cDNA and its relationship with a loss of *rbcS* in potato (*Solanum tuberosum* L.) plants. *Plant Mol Biol* **28**: 93–103
- Schraudner M, Moeder W, Wiese C, van Camp W, Inzé D, Langebartels C, Sandermann H Jr (1998) Ozone-induced oxidative burst in the ozone biomonitor plant, tobacco Bel W3. *Plant J* **16**: 235–245
- Serek M, Tamari G, Sisler EC, Borochov A (1995) Inhibition of ethylene-induced cellular senescence symptoms by 1-methylcyclopropane, a new inhibitor of ethylene action. *Physiol Plant* **94**: 229–232
- Shirasu K, Nakajima H, Rajasekhar VK, Dixon RA, Lamb C (1997) Salicylic acid potentiates an agonist-dependent gain control that amplifies pathogen signals in the activation of defense mechanisms. *Plant Cell* **9**: 261–270
- Siefert F, Kwiatkowski J, Sarkar S, Grossmann K (1995) Changes in endogenous cyanide and 1-aminocyclopropane-1-carboxylic acid levels during the hypersensitive response of tobacco mosaic virus-infected tobacco leaves. *Plant Growth Regul* **17**: 109–113
- Simons BH, Millenaar FF, Mulder L, Van Loon LC, Lambers H (1999) Enhanced expression and activation of the alternative oxidase during infection of *Arabidopsis* with *Pseudomonas syringae* pv. *tomato*. *Plant Physiol* **120**: 529–538
- Staskawicz BJ, Ausubel FM, Baker BJ, Ellis JG, Jones JDG (1995) Molecular genetics of plant disease resistance. *Science* **268**: 661–667
- Thomma BPHJ, Eggermont K, Penninckx IAMA, Mauch-Mani B, Vogelsang R, Cammue BPA, Broekaert WF (1998) Separate jasmonate-dependent and salicylate-dependent defense-response pathways in *Arabidopsis* are essential for resistance to distinct microbial pathogens. *Proc Natl Acad Sci USA* **95**: 15107–15111
- Tuomainen J, Betz C, Kangasjärvi J, Ernst D, Yin ZH, Langebartels C, Sandermann H Jr (1997) Ozone induction of ethylene emission in tomato plants: regulation by differential transcript accumulation for the biosynthetic enzymes. *Plant J* **12**: 1151–1162
- Vahala J, Keinänen M, Schutzendubel A, Polle A, Kangasjärvi J (2003) Differential effects of elevated ozone on two hybrid aspen genotypes predisposed to chronic ozone fumigation: role of ethylene and salicylic acid. *Plant Physiol* **132**: XXX–XXX
- Vahala J, Schlagghauser CD, Pell EJ (1998) Induction of an ACC synthase cDNA by ozone in light-grown *Arabidopsis thaliana* leaves. *Physiol Plant* **103**: 45–50

- Wenzel AA, Schlautmann H, Jones CA, Küppers K, Mehlhorn H** (1995) Aminoethoxyvinylglycine, cobalt and ascorbic acid all reduce ozone toxicity in mung beans by inhibition of ethylene biosynthesis. *Physiol Plant* **93**: 286–290
- Wilkinson JQ, Lanahan MB, Clark DG, Bleecker AB, Chang C, Meyerowitz EM, Klee HJ** (1997) A dominant mutant receptor from *Arabidopsis* confers ethylene insensitivity in heterologous plants. *Nat Biotechnol* **15**: 444–447
- Wohlgemuth H, Mittelstrass K, Kschieschan S, Bender J, Weigel HJ, Overmyer K, Kangasjärvi J, Langebartels C, Sandermann H** (2002) Activation of an oxidative burst is a general feature of sensitive plants exposed to the air pollutant ozone. *Plant Cell Environ* **25**: 717–726
- Wurtele ES, Nikolau BJ, Conn EE** (1985) Subcellular and developmental distribution of  $\beta$ -cyanoalanine synthase in barley (*Hordeum vulgare*) leaves. *Plant Physiol* **78**: 285–290
- Yip W-K, Yang SF** (1988) Cyanide metabolism in relation to ethylene production in plant tissues. *Plant Physiol* **88**: 473–476
- Ziegler J, Keinänen M, Baldwin IT** (2001) Herbivore-induced allene oxide synthase transcripts and jasmonic acid in *Nicotiana attenuata*. *Phytochemistry* **58**: 729–738